

AD-A088 37\*

TENNESSEE UNIV KNOXVILLE DEPT OF MATHEMATICS  
A ZERO-ONE LAW FOR A CLASS OF MEASURES ON GROUPS.(U)  
SEP 79 B RAJPUT, D LOUIE, A TORTRAT

F/6 12/1

N00014-78-C-0468

UNCLASSIFIED

TR-7905

NL

1 of 1  
AD-A088 37\*



END  
DATE  
FILMED  
9-80  
DTIC


LEVEL II

DEPARTMENT OF MATHEMATICS  
The University of Tennessee, Knoxville


12

54

AD A088374



DTIC  
ELECTE  
AUG 26 1980



DDC FILE COPY

This document has been approved  
for public release and sale; its  
distribution is unlimited.

80 8 21 059

12

A ZERO-ONE LAW FOR A CLASS OF MEASURES ON GROUPS

by

Donald Louie and Balram S. Rajput  
The University of Tennessee, Knoxville, U.S.A.

and

Albert Tortrat  
The University of Paris, France

DTIC  
ELECTE  
AUG 26 1980  
S D C

This document has been approved  
for public release and sale; its  
distribution is unlimited.

# A ZERO-ONE LAW FOR A CLASS OF MEASURES ON GROUPS

by

Donald Louie  
The University of  
Tennessee  
Knoxville, Tennessee

Balram S. Rajput\*  
The University of  
Tennessee  
Knoxville, Tennessee

Albert Tortrat  
The University of Paris,  
France

## ABSTRACT

Let  $(E, \mathcal{B})$  be a measurable (non-abelian) group,  $\alpha$  a measurable homomorphism on  $E$  and  $\mu$  a law on  $(E, \mathcal{B})$  satisfying  $\mu = \mu_\alpha \nu_\alpha$  (with  $\mu_\alpha = \mu \circ \alpha^{-1}$ ), for some law  $\nu_\alpha$  on  $(E, \mathcal{B})$ . Under some additional conditions on  $\mu, \nu_\alpha$  and  $\alpha$ , it is shown that, for every proper normal subgroup  $G$  of  $E$  and  $a$  in  $E$ , if  $Ga \in \mathcal{B}_\mu^*$ , the  $\mu$ -completion of  $\mathcal{B}$ , then  $\mu(Ga) = 0$  or  $1$ . As a corollary, it is shown that this result yields all previously known 0-1 laws for stable, semistable, and quasistable laws on linear spaces as well as new 0-1 laws for other classes of infinitely divisible laws on linear spaces.

\* The research of this author is partially supported by the Office of the Naval Research under contract No. N0014-78-C-0468.

Classification For	<input checked="" type="checkbox"/> CONFIDENTIAL <input type="checkbox"/> SECRET <input type="checkbox"/> UNCLASSIFIED
DDC TAB	
Unannounced	
Justification	
By	
Distribution	
Availability Codes	
Available for	
Dist Special	

The main purpose of this paper is to prove the

Theorem: We suppose that  $(E, \mathcal{B})$  is a measurable (non-abelian) group (i.e.  $E$  is a non-abelian group,  $\mathcal{B}$  is a  $\sigma$ -algebra of subsets of  $E$  such that the map  $x \times x' \rightarrow xx^{-1}$  is measurable relative to the  $\sigma$ -algebras  $\mathcal{B} \times \mathcal{B}$  and  $\mathcal{B}$ ) or that  $(E, \mathcal{B})$  is a Hausdorff topological group with  $\mathcal{B}$  its Borel  $\sigma$ -algebra. In the latter case all laws considered are assumed to be  $\tau$ -regular. Let  $\alpha$  be a measurable homomorphism from  $E$  into  $E$  (i.e. a measurable (relative to  $\mathcal{B}$  and  $\mathcal{B}$ ) map preserving the group operation) and  $\mu$  a law on  $\mathcal{B}$ ; and let  $\mu_\alpha$  and  $\mathcal{B}_\mu$  denote, respectively, the image of  $\mu$  under  $\alpha$  and the  $\mu$ -completion of  $\mathcal{B}$ . We assume that  $\mu_\alpha$  is a left factor of  $\mu$  (on  $\mathcal{B}$  and hence on  $\mathcal{B}_\mu$ ) and an  $n$ -th root of  $\mu_\alpha$  exists which divides the cofactor  $\nu_\alpha$ :

$$(1) \quad \mu = \mu_\alpha \nu_\alpha, \quad (1') \quad \nu_\alpha = \mu^{1/n} \nu'_\alpha \quad \text{or} \quad \nu_\alpha = \nu'_\alpha \mu^{1/n}.$$

Let  $G$  be a normal subgroup of  $E$  and assume that

$$(2) \quad \alpha(G) \subseteq G, \quad \alpha(G^c) \subseteq G^c, \quad (2')^* \quad \alpha^n I^{-1}(G^c) \subseteq G^c,$$

for all  $n = 1, 2, \dots$ , where  $\alpha^n$  denotes the  $n$ -th iteration of  $\alpha$  and  $G^c$  denotes the complement of  $G$ . Then, for any  $a \in E$ , we have

$$(3) \quad Ga \in \mathcal{B}_\mu \Rightarrow \mu(Ga) = 0 \text{ or } 1.$$

---


$$(*) \quad (\alpha^n I^{-1})(x) = \alpha^n(x) x^{-1}, \text{ for } x \in E.$$

If  $G = \{e\}^+$  and  $\{a\} \in \mathcal{B}_\mu$ , for all  $a \in E$ , or if  $G$  is closed, then we can, in the topological case, replace (1') and (2') by

$$(1'') \quad \lim_n \alpha^n(x) = e, \text{ for all } x \in E.$$

Proof. Assume  $\mu(Ga) > 0$ , we will show  $\mu(Ga) = 1$ . Let  $G_0$  denote the group of right cosets  $Gx$ . Condition (2) assures us that  $\alpha$  induces in  $G_0$  an injective homomorphism. We denote the coset  $Gc$  by  $\dot{c}$ ; if  $\dot{c} \in \mathcal{B}_\mu$ , then, from Fubini's theorem and (1), we have (with  $\nu \equiv \nu_\alpha$ )

$$(4) \quad \mu(\dot{c}) = \int_E \mu\{\alpha^{-1}(\dot{c}x^{-1})\} \nu(dx).$$

In (4),  $\alpha^{-1}(\dot{c}x^{-1}) \in \mathcal{B}_\mu$  and  $g(x) = \mu\{\alpha^{-1}(\dot{c}x^{-1})\}$  is  $\mathcal{B}$ -measurable off a set  $N \in \mathcal{B}$  with  $\nu(N) = 0$ .

Let  $A = \{\dot{c}_k \equiv Gc_k : k = 1, \dots, K\}$  form the finite set of all distinct  $\mu$ -measurable right cosets of  $G$  such that  $\mu(\dot{c}_k)$  is maximum (among all  $\mu$ -measurable right cosets). We will show  $K = 1$  by showing that  $K \geq 2$  leads to a contradiction. From (4) and the maximality of  $\mu(\dot{c}_k)$ , one can choose distinct cosets  $\dot{b}_i^{-1}$ ,  $i = 1, \dots, I$ ,  $I \leq K$ , such that  $\nu(\cup \dot{b}_i^{-1}) = 1$  and  $\alpha^{-1}(\dot{c}_k \dot{b}_i) = \dot{c}_{ki} \in A$ , with  $\dot{c}_{ki}$  distinct (for fixed  $i$ ) (we don't know that  $\dot{b}_i^{-1}$  is  $\nu$ -measurable). Denote by  $\Delta$  the set of symmetric (distinct)

---

<sup>†</sup> This case, when  $\mu$  is infinitely divisible and  $E$  is a topological space, comes up in the study of atoms of these laws [1]; the results of [1] trivially extend from the cylindrical  $\sigma$ -algebra to  $\mathcal{B}$  when  $\mu$  is  $\tau$ -regular.

elements  $\dot{c}_k \dot{c}_{k'}^{-1}$ ,  $k \neq k'$ , say  $\Delta \equiv \{\delta_\ell; \ell = 1, \dots, L\}$  with  $1 \leq L$  (as  $K \geq 2$ ). Then since  $\dot{c}_k \neq \dot{c}_{k'}$  implies  $\dot{c}_{ki} \neq \dot{c}_{k'i}$  and  $\alpha^{-1}(\dot{c}_k \dot{c}_{k'}^{-1}) = \alpha^{-1}(\dot{c}_k b_i b_i^{-1} \dot{c}_{k'}^{-1}) = \dot{c}_{ki} \dot{c}_{k'i}^{-1}$ , it follows that

$$\alpha^{-1}(\delta_\ell) = \delta_\ell, \in \Delta.$$

Thus  $\alpha^{-1}$  is a permutation in  $\Delta$ ; hence  $\alpha^n$  has a fixed point, say  $\dot{b}$ , for some  $n \leq L$ . In  $E$  this is written:  $\alpha^n(b) = gb$ ,  $g \in G$ ; thus  $\alpha^{n-1}(b) \in G$ . Then (2') implies  $b \in G$ ; thus  $\dot{b} = \epsilon$ , the unit element of  $G_0$ , but  $\epsilon \notin \Delta$ . This is a contradiction; hence  $K = I = 1$ . Hence  $v(\dot{c}_1) = 1$ ; but then (1') and  $\mu(\dot{a}) > 0$  imply  $\mu(\dot{a}) = 1$ . In the case that  $G = \{e\}$  and in order to replace (1') by (1''), we make use of the following:

Lemma. Let  $G_0$  be a Hausdorff topological group with unit element  $\epsilon$   $\mu$  a law (of a random variable  $X$ ) on the Borel  $\sigma$ -algebra  $\mathcal{B}_0$  of  $G_0$ . We assume  $\mu$  is  $\tau$ -regular and has nonempty support (if  $G_0$  is metrizable) and  $\alpha$  is as given above from  $E$  into  $E$ . Then the conditions

$$(5) \quad \alpha(X) \stackrel{\text{law}}{=} Xb \quad \text{and} \quad (1'') \quad \alpha^n(x) \xrightarrow{n \rightarrow \infty} \epsilon, \quad \text{for all } x,$$

imply that  $\mu$  is improper:  $X = a$  a.s. with  $\alpha(a) = ab$ .

Proof. Let  $V$  be a symmetric neighborhood of  $\epsilon$  and  $\mu(Va) = \eta > 0$ . Then (1'') implies that

$$\{x : \alpha^n(x) \in V, \text{ all } n \geq m\} \uparrow G_0 \text{ when } m \uparrow \infty.$$

On the otherhand (5) gives

$$\alpha^n(X) \stackrel{\text{law}}{=} X b_n, \quad \text{with } b_n = \alpha^{n-1}(b) \dots \alpha(b)b.$$

Thus, we have  $P\{\alpha^n(X)b_n^{-1} = X \in Vb_n^{-1}\} = P\{\alpha^n(X) \in V\} + \dots$ .

Hence,  $Vb_n^{-1}$  intersects  $Va$  for all  $n \geq n_0$  sufficiently large; thus  $b_n^{-1} \in V^2a$  and  $P\{X \in V^3a\} \geq P\{X \in Vb_n^{-1}\} + 1$ . Hence the  $\tau$ -regularity of  $\mu$  or the weaker hypothesis that its support is nonempty (if  $G_0$  is metrisable) assures  $X = a$  a.s. That  $\alpha(a) = ab$  is clearly true.

Remark. Even if  $E$  is a Hausdorff topological group,  $G$  nonclosed (which is the interesting case) does not assure that  $G_0$  is such a group. Then it is very possible that, with the atom  $\dot{a}$  of  $G_0$  removed, there remains a single atom, for the law  $\mu$  is zero on the points of  $G_0$ . In this case the relation  $\alpha(X) \stackrel{\text{law}}{=} Xb$  is possible\*, and yet  $\mu(\dot{a}) = 1$  is not necessarily true (: the  $G$ -cylinders of  $E$ , belonging to  $\mathcal{B}_\mu$ , except  $Ga$ , reduce to the countable union of classes which have  $\mu$ -measure 0 and to the complements of these unions which have  $\mu$ -measure 1, and are equivalent to the above mentioned atom). Thus hypothesis (1'), which appears artificial except in the case when  $\mu$  is infinitely divisible, seems indispensable.

Corollary. Let  $E$  be a measurable ( $x \times x' \rightarrow x + x'$  and  $t \times x \rightarrow tx, t$  real, are measurable) or a Hausdorff topological vector space. Let  $0 < \alpha < 1$ ,

---

\*  $G_0 \setminus \{\dot{a}\}$  can also, for example, consist of two equal atoms which can be exchanged under the action of  $\alpha$  or two unequal (and thus) invariant atoms.

By hypothesis these atoms are not reduced to the points of  $G_0$  with zero  $\mu$  measure.



$\alpha(\cdot)$  the map  $\alpha(x) = \alpha \cdot x$ , and  $\ell(\alpha)$  the field generated in  $R$  by  $\alpha$ . Let  $G$  be a additive subgroup of  $E$ . We retain for  $\mu$  the conditions (1) and (1') (and only (1) if  $E$  is a Hausdorff topological vector space and  $G = \{0\}$ ). Then  $G' + a \in \mathcal{B}_\mu$  (with  $G' = \ell(\alpha)G$ ) implies

$$(6) \quad \mu(G' + a) = 0 \text{ or } 1.$$

Thus,  $G \in \mathcal{B}$  also implies (6).

Proof.  $G'$  satisfies (2) and (2').

This applies to the quasi-stable, semi-stable [5], and stable laws [4] without passing to symmetrisation or to centering of laws, but we can not deduce  $G \in \mathcal{B}_\mu \Rightarrow \mu(G+a) = 0 \text{ or } 1$ , a property known for a centered Gaussian law (: stable of exponent 2). In fact for  $\alpha$  rational the condition (2') along with  $\alpha G = G$  (note that for any bijective map  $\alpha$  on  $E$ , not necessarily a homomorphism, the condition  $\alpha(G) = G$  is equivalent to (2)) imply that  $G$  is a vector space over the field  $\mathbb{Q}$  of rationals and thus the theorem applies only to certain groups.

Other examples: Let  $E$  be a measurable or a Hausdorff topological vector space,  $\mu = P(M)$  be a Poisson type infinitely divisible law on  $E$ , and  $\alpha$  a 1-1 bounded linear operator on  $E$ . The condition

$$(7) \quad \alpha(M) \equiv M_\alpha \leq \frac{n-1}{n} M$$

(for large enough  $n$ ) implies that  $\mu_\alpha = P(M_\alpha)$  satisfies (1) and (1'). Then every  $G + a$  satisfying (2) and (2') is of  $\mu$ -measure 0 or 1 (but the condition (2') is difficult to interpret and verify). Thus, when

Then every  $G + a$  satisfying (2) and (2') is of  $\mu$ -measure 0 or 1 (but the condition (2') is difficult to interpret and verify). Thus, when  $\alpha\{\cdot\}$  is multiplication by a number  $\alpha \in (0,1)$ , one obtains 0-1 laws for a certain class of self-decomposable laws (see [2]) as well as for many others (e.g. Radon  $\alpha$ -decomposable laws  $P(M)$  with  $M$  satisfying (7)). An analogous remark applies (with  $\alpha$  1-1 linear and  $\alpha$  and  $\alpha^{-1}$  bounded) to the laws of class  $L$  studied by Urbanik in [3].

#### REFERENCES

- [1] Albert Tortrat, Atomes et lois indéfiniment divisibles dan un espace vectoriel, Ann. Inst. H. Poincaré XIV, 3, 1978, 345-349.
- [2] Dong M. Chung, Characterization of self decomposable probability measures on LCTVS, J. Korean Math. Soc., 15-2 (1979), 139-148.
- [3] K. Urbanik, Lévy's probability measures on Banach spaces, Studia Mathematica, LXIII (1978), 283-308.
- [4] R.M. Dudley and M. Kantor, Zero-one laws for stable measures, Proc. Amer. Math. Soc., 45 (1974), 245-252.
- [5] D. Louie, Balram S. Rajput and Albert Tortrat, A zero-one dichotomy theorem for  $r$ -semistable laws on infinite dimensional linear spaces, Sankhyā, (1979), (in press).

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report # 7905	2. GOVT ACCESSION NO. AD-A088 374	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A 0-1 Law for A Class Of Measures On Groups.		5. TYPE OF REPORT & PERIOD COVERED
6. PERFORMING ORG. REPORT NUMBER		7. AUTHOR(s) Balram/Rajput, Donald/Louie, Albert/Tortrat, Univ. of Paris, France
8. CONTRACT OR GRANT NUMBER(s) N00014-78-C-0468		9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Department, University of Tennessee, Knoxville, TN 37916
10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-042-400		11. CONTROLLING OFFICE NAME AND ADDRESS Statistics and Probability Program, Office of Naval Research, Arlington, Va 22217
12. REPORT DATE September, 1979		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (12) 11		15. SECURITY CLASS. (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Semi-stable, stable, Group zero-one laws		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let $(E, \mathcal{B})$ be a measurable (non-abelian) group, $\alpha$ a measurable 1-1 homomorphism on $E$ and $\mu$ a law on $(E, \mathcal{B})$ satisfying $\mu = \mu_\alpha \nu_\alpha$ (with $\mu_\alpha = \mu \circ \alpha^{-1}$ ), for some law $\nu_\alpha$ on $(E, \mathcal{B})$ . Under some additional conditions on $\mu, \nu_\alpha$ and $\alpha$ , it is shown that, for every proper normal subgroup $G$ of $E$ and $a$ in $E$ , if $Ga \in \mathcal{B}_\mu$ , the $\mu$ -completion of $\mathcal{B}$ , then $\mu(Ga) = 0$ or 1. As a corollary		

DD FORM 1473 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE  
S N 3132-LF-014-5601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

411 894

JB

Abstract(continued)

it is shown that this result yields all previously known 0-1 laws for stable, semistable, and quasistable laws on linear spaces as well as new 0-1 laws for other classes of infinitely divisible laws on linear spaces.